

**This product helps to protect chlorine in pool water from dissipation caused by sunlight. Use if intense sunlight causes difficulty in maintaining chlorine residual.**

1. Clean and backwash the filter.
2. Adjust chlorine level to 1-3 ppm with a shock treatment.
3. Measure the stabilizer (cyanuric acid) concentration with an appropriate test kit. The Ideal range is 20-25 ppm. Do not exceed 100 ppm as chlorine effectiveness will be reduced.
4. Determine dosage of this product from the table below .
5. With circulation pump on, add this product to the skimmer . Leave the circulation pump running for 48 hours.
6. Check the stabilizer level at least twice each year and add as needed.

Pool Volume in Gallons	To Raise 10 ppm	To Raise 20 ppm	To Raise 25 ppm
10,000	1	2	2.5
20,000	2	4	5
40,000	4	8	10

**Sold by:**  
**Robelle Industries, Inc.**  
**131 Tosca Drive**  
**Stoughton, MA 02072**



# CHLORINE STABILIZER and Conditioner

*everything for your swimming pool*

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**See First Aid Statement and additional precautions on back panel. Read all instructions carefully before use.**

**Isocyanuric Acid ..... 100%**

ITEM NO.  
L9007BA

**Storage:** Keep product dry in tightly closed container when not in use. Store in a cool, dry, well ventilated area. Keep away from heat or open flame. **Disposal:** Do not reuse empty container. Rinse empty container. Offer for recycling if available or place in trash collection. **If Partly Filled:** Call your local solid waste agency or 1-800-CLEANUP for instructions. Never place unused product down any indoor or outdoor drain.

**Caution: Eye irritant. Avoid contact with eyes. Do not mix with any other chemicals. Contamination may cause fire. Keep out of reach of children. Contains cyanuric acid.**

### If Swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

- Hold eye open and gently rinse with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

**Have the product container or label with you when calling a poison control center or doctor or going for treatment.**